

CORRECTION

Open Access



Correction to: Phytochemical studies, antiangiogenic, antiinflammatory and antioxidant activities of *Scyphocephalum ochocoa* Warb. (Myristicaceae), medicinal plant from Gabon

Rick-Leonid Ngoua-Meye-Misso^{1,2,5*}, Jean De La Croix Ndong³, Cédric Sima-Obiang^{1,2}, Joseph Privat Ondo^{1,2}, Guy Roger Ndong-Atome^{1,2}, Felix Ovono Abessolo⁴ and Louis-Clément Obame-Engonga^{1,2}

Correction

Following the publication of this article [1], the authors noticed that Figs. 1, 2 and 3 were in the incorrect order and thus had incorrect captions. The images that were incorrectly published as Figs. 1, 2 and 3 should have been published as Figs. 2, 3 and 1 respectively.

The correct versions of Figs. 1, 2 and 3 with captions have been included in this Correction.

The original article has been corrected.

Author details

¹Laboratory of Research in Biochemistry (LAREBIO), University of Sciences and Technology of Masuku, Franceville, Gabon. ²Laboratory of Natural Substances and Organometallic Synthesis, University of Sciences and Technology of Masuku, Franceville, Gabon. ³Department of Otolaryngology-Head and Neck Surgery, Ear, Nose and Throat Laboratory, Emory School of Medicine, Atlanta, USA. ⁴Laboratory of Biochemistry, Joint Unit of Biomedical Research, University of Health Sciences, Libreville, Gabon. ⁵Department of Chemistry and Biochemistry, Faculty of Science, University of Science and Technology of Masuku, Franceville, Gabon.

Received: 14 June 2018 Accepted: 18 June 2018

Published online: 02 July 2018

Reference

1. Ngoua-Meye-Misso et al. *Clinical Phytoscience*. 2018;4:15. <https://doi.org/10.1186/s40816-018-0075-x>.

* Correspondence: ngouamartel@gmail.com; ngouamartel@univ-masuku.ga

¹Laboratory of Research in Biochemistry (LAREBIO), University of Sciences and Technology of Masuku, Franceville, Gabon

²Laboratory of Natural Substances and Organometallic Synthesis, University of Sciences and Technology of Masuku, Franceville, Gabon



Fig. 1 Shrub and trunk of *S. ochocoa*. Photos taken at Mitzic, Woleu-Ntem (Northern of Gabon) by Ngoua-Meye-Misso with digital camera Cannon 16 M pixel made in China




Fig. 2 Inhibitory effects of water extracts of plants on angiogenesis. The CAM of a 8 days old chick embryo was separately exposed to PBS (control). Extracts were introduced on top of the CAMs. After 48 h of incubation, the CAM tissue directly beneath each filter disk was resected, and digital images of the CAM sections were captured




Fig. 3 Haemoglobin content of CAM sections detected by Drabkin's assay. The hemoglobin content in the CAM sections was determined by Drabkin method. As in the case of macroscopic observation of CAM that showed that the number of blood vessel branches was concentration dependent, so the present figure shows a hemoglobin content dependent on the concentration of *S. Ochococa* water extract and sorafenib