

ORIGINAL CONTRIBUTION

Open Access



Screening of phytochemical and pharmacological activities of *Syzygium caryophyllatum* (L.) Alston

Stalin N and Sudhakar Swamy P*

Abstract

Background: The systematic screening of plant species with the purpose of discovering new bioactive compounds is prerequisite for any bioprospecting study. Therefore the present study was carried out to assess the phytochemical content and the evaluation of in vitro antioxidant, antibacterial, larvicidal and antidiabetic activities of the methanol extracts of *S. caryophyllatum* bark, leaves, fruit pulp and seeds.

Methods: The quantitative estimation of total phenol, flavonoid and tannin content of the extracts of *S. caryophyllatum* bark, leaf, fruit pulp and seeds were analyzed by using standard methods. The present study also conducted to screen antioxidant (DPPH, ABTS assays), antimicrobial, antidiabetic (in vitro α -amylase and α -glucosidase inhibitory assays) and larvicidal (against fourth instars larvae of *Aedes aegypti* and *Culex quinquefasciatus*) activities of methanol extract of *S. caryophyllatum*.

Results: The results of quantitative phytochemical analysis revealed the presence of maximum amount of phytoconstituents such as phenol, flavonoids and tannins in the leaf, bark and seed extract when compared to fruit pulp extracts. Free radical scavenging activity indicated that methanol bark, leaves, fruit and seeds extracts has significant free radical scavenging ability on DPPH with percentage inhibition of 88.15%, 81.31%, 75.24% and 83.36% respectively. The in vitro α -amylase and α -glucosidase inhibitory studies of the methanol crude extracts of four different plant parts of *S. caryophyllatum* (Bark, leaf fruit pulp and seed) showed good inhibitory activities in concentration dependent manner. The maximum percentage inhibitory activity of 78.03% was showed at concentration of 500 μ g/ml seed extracts followed by bark (78.03%), leaf (69.4%) and fruit pulp (56.9%) at the same concentration. While the percentage inhibitory activity of four extracts also showed potent inhibition of α -glucosidase; maximum inhibition exhibited at the concentration of 100 μ g/ml by bark extract (80.9%) compared with other extracts, leaf (78.2%), seed (77.59%) and fruit pulp (63.35%). The leaf essential oil and four extracts showed significant mortality against fourth instars larvae of *Ae. aegypti* and *Cx. quinquefasciatus* respectively.

Conclusion: Thus the present study suggests that *S. caryophyllatum* plant parts can be used as natural antioxidant source to prevent diseases associated with free radicals. Also, this plant can be a good source for further purification studies for isolation and characterization of compounds related to these antioxidants, antidiabetic and antibacterial activities.

Keywords: *Syzygium caryophyllatum*, DPPH, ABTS, Larvicidal, Free radicals

* Correspondence: swamypsplantscience@gmail.com
Department of Plant Science, School of Biological Sciences, Madurai Kamaraj University, Madurai, Tamil Nadu -21, India

Background

Nature has been a resource of medicinal treatments for thousands of years, and plant derived products being continued to play a crucial role in primary health care of approximately 80% of the world's population [1]. Plant based drugs can be derived from any part of the plant like leaves, bark, flowers, fruits, root etc. i.e., any part of the plant may contain active components. The systematic screening of plant species with the purpose of discovering new bioactive compounds is prerequisite for any bioprospecting study. The fruits from wild edible plants possess significant amount of vitamins (β -carotene, vitamins C and E), flavonoids and other polyphenolic compounds having high free-radical scavenging activity [2]. An epidemiological study provides convincing evidence of the beneficial role of fruits and vegetables in the diet for the maintenance of health and prevention of diseases [3]. Polyphenols represent a diverse class of compounds including flavonoids (i.e. flavones, flavonols, flavanones, flavononols, flavan-3-ols and chalcones), tannins and phenolic acids [4]. These polyphenols are products of secondary metabolism in plants, and their important role of chelating transitional metals and scavenging free radicals activity are mainly due to their redox properties and chemical structures [5]. Antioxidants are vital substances which possess the ability to protect the body from tissue damage caused by free radical induced oxidative stress [6]. In addition to common antioxidants like vitamin C, vitamin E and carotenoids, phenolic constituents such as flavonoids, tannins and anthocyanins present in food of plant origin are also potential antioxidants [7–9].

Plant phenolics present in the fruit and vegetables have received considerable attention because of their potent antioxidant property. Plants possess more phenolic content show better antioxidant activity this is due to the direct correlation between total phenolic content and antioxidant activity [10, 11]. Natural products of higher plants may possess a new source of antimicrobial agents with possibly novel mechanism of action [12, 13]. They are effective in the treatment of infectious diseases while simultaneously mitigating many of the side effects that are often associated with synthetic antimicrobials [14].

Mosquitoes are the most important single group of insects in terms of public health importance, which are responsible for spreading a number of severe diseases, such as malaria, filariasis, dengue, Japanese encephalitis, etc. *Culex quinquefasciatus* is a major contributor for transmitting vector borne diseases like fevers and filariasis to human beings and animals. More than 40 million people affected globally from Elephantiasis and are badly debilitated and disfigured from this disease [15]. *Aedes* mosquitoes on the other hand are vectors of important

viral diseases of man such as the zika virus, dengue and yellow fevers. *Ae. aegypti* is reported to be the principal vector of dengue fever in tropical countries, a disease considered a serious public health problem world over, especially in areas where the environmental conditions for the development of *Aedes* mosquitoes abound. Totally, 2.35 million people suffering from dengue was reported in America alone (as per 2013 data) of which 7687 cases were rigorous [16]. According to Directorate of National Vector Borne Disease Control Programme (NVBDCP), till July 30, 2016, the total dengue cases in India were 16,870 while for the same period in 2017 they numbered 28,702. Recently, the infections have been raised due to an increased urbanization, trade, and travel. No effective drug or vaccine is available so far. Frequent uses of these synthetic insecticides create an increase resistance among the mosquitoes and to cause adverse impacts on non-target organisms or humans. With this scenario, biological control especially plant based insecticide has proven as better in controlling insect vectors.

Plant extracts have long been used for the treatment of diabetes in various traditional systems of medicine and are now accepted as an alternative for diabetic treatment. Type II diabetes mellitus is a chronic endocrine metabolic disorder that affects the carbohydrates, proteins and fat metabolism. Postprandial increase of blood glucose is the major problem in diabetic patients. Inhibitors of α -amylase and α -glucosidase enzymes delay the breaking down of carbohydrates in the small intestine and decrease the postprandial blood glucose levels. Inhibitors of α -amylase and α -glucosidase isolated from medicinal plants to serve as an alternative drug with increased potency and lesser adverse effects than existing synthetic drugs.

Syzygium caryophyllatum (L.) Alston., an endangered evergreen tree belongs to the family Myrtaceae, commonly known as Wild black plum. This tree is native to Sri Lanka and India; in India the distribution is restricted to the forests of Western Ghats regions. *S. caryophyllatum* fruits are edible, sweet and astringent in taste. The seeds and bark were dried and its decoction was used in the ailment of diabetes mellitus [17]. The leaf and bark extracts of this plant are well known for its antibacterial and antioxidant efficacy [18]. Root extract using different solvent system exhibited promising amount of phenol, flavonoids, tannins and vitamin C [19]. The bark extract of this plant was used in veterinary medicine for the treatment of tympanitis in cattles [20]. Tribal people consider this plant as a boon of nature and its fruits and seeds were consumed by Paniya tribal community of Waynad district, Kerala, India [21]. As the plant is used for medicinal purposes, the investigation of chemical constituents in the plant parts

and its biological activities may provide valuable information. There is a detailed bioprospecting studies by using different organic solvent system with different plant parts have not been studied previously on *S. caryophyllatum*. Therefore the present study was carried out (i) to analyse the leaves essential oil and its antibacterial and larvicidal activities (ii) to assess the total phenols, flavonoids and tannins content; (iii) to evaluate the in vitro antioxidant, antibacterial, larvicidal and antidiabetic activities of the methanol extracts of *S. caryophyllatum* bark, leaves, fruit pulp and seeds.

Methods

Plant material and sample preparations

S. caryophyllatum plant samples (Leaves, bark and fruits) were collected from Kannur district of Kerala, India during 2012 and voucher specimen (PLS2011007) was prepared, identified and deposited at the Department of Plant Science herbarium, Madurai Kamaraj University. The bark, leaves, fruit pulp and seeds were separately dried at below 40 °C. Powdered sample of *S. caryophyllatum* bark, leaves, fruit pulp and seeds was sequentially extracted with hexane, ethyl acetate, methanol and water in their increasing polarity by sonicator. The extracts were filtered through Whatman No.1 filter paper and the solvents were evaporated under reduced pressure. Dried extracts obtained were stored in 4 °C until further analysis.

Estimation of total phenol content

The amount of total phenol in plant extracts was determined with the Folin–Ciocalteu reagent using the method given by Lister and Wilson [22]. 100 µl of appropriate dilutions of extracts were dissolved in 500 µl (1/10 dilution) of the Folin–Ciocalteu reagent and 1000 µl of distilled water. The solutions were mixed and incubated at room temperature for 1 min. After 1 min, 1500 µl of 20% sodium carbonate (Na₂CO₃) solution was added to the mixture and then incubated for 2 h in the dark at room temperature. A standard curve was prepared by using gallic acid as a standard. Different concentrations of gallic acid were prepared in 80% of methanol. The absorbances of all samples were measured at 760 nm using a UV-Vis spectrophotometer (Model. U.2800, Hitachi) and the total phenol contents were expressed as mg/g gallic acid equivalent (GAE) using the equation obtained from the calibration curve using the following eqs.

$$Y = b \times x + c \quad (1)$$

Where, Y-absorbance of extract; c-intercept; b-slope of the calibration curve; x- concentration of extract.

$$X = Y - c/m \quad (2)$$

$$C = cV/m \quad (3)$$

Where, C-Total phenolic content; c-concentration of gallic acid obtained from calibration curve in mg/ml; V-volume of extract in ml; m- mass of extract in gram.

Estimation of total flavonoid content

Estimation of total flavonoid content in the dried extracts was determined spectrophotometrically following the method given by Quettier-Deleu et al. [23]. This method is based on the formation of a complex flavonoid–aluminium, having the absorbivity maximum at 430 nm. 1 ml of diluted sample was separately mixed with 1 ml of aluminum chloride solution (2% w/v). After incubation at room temperature for 15 min, the absorbance of the reaction mixture was measured at 430 nm. Rutin was used to make the calibration curve. The flavonoid content was expressed in mg per g of rutin equivalent (RE using the equation obtained from the calibration curve using the following eqs.

$$Y = bx + c \quad (1)$$

Where, Y-absorbance of extract; c-intercept; b-slope of the calibration curve; x- concentration of extract.

$$X = Y - c/m \quad (2)$$

$$C = cV/m \quad (3)$$

Where, C-Total flavonoid content; c-concentration of rutin obtained from calibration curve in mg/ml; V-volume of extract in ml; m- mass of extract in gram.

Estimation of tannin content

The amount of tannin content was determined by modified Prussian blue method [24]. This method is based on the mechanism that the phenols reduce potassium ferricyanide to produce ferrous ions; these ferrous ions in turn react with ferric chloride in the presence of dilute HCl to form a Prussian blue coloured complex, which can be measured spectrophotometrically at 700 nm wavelength. 100 µl of appropriate dilutions of extracts were dissolved in distilled water to make up the volume of 7 ml. The sample solution was mixed with 1 ml potassium ferricyanide followed by the addition of 1 ml ferric chloride. The absorbance of samples was measured at 700 nm. Tannic acid was used to make a calibration curve. The amount of tannin in plant extracts was calculated by using the equation obtained from the calibration curve using the following eqs.

$$Y = bx + c \quad (1)$$

Where, Y-absorbance of extract; c-intercept; b-slope of the calibration curve; x- concentration of extract.

$$X = Y - c/m \quad (2)$$

$$C = cV/m \quad (3)$$

Where, C-Total Tannin content; c-concentration of Tannic acid obtained from calibration curve in mg/ml; V-volume of extract in ml; m- mass of extract in gram.

In vitro free radical scavenging assays

Free-radical scavenging ability (DPPH-assay)

The scavenging ability of leaf essential oil, bark, leaves, fruit pulp and seed extracts on DPPH free- radicals was estimated according to the method of Shimada et al. (1992) [25]. This method depends on the reduction of purple 1,1-diphenyl-2-picrylhydrazyl (DPPH) to a yellow coloured diphenyl picrylhydrazine and the remaining DPPH which showed maximum absorption at 517 nm was measured. About 2 ml of various concentrations of test samples and ascorbic acid and gallic acid standards (2.5–20 µg/ml) were mixed with 0.5 mL of 1 mM DPPH in methanol. An equal amount of methanol and DPPH served as control. The mixture was shaken vigorously and then incubated for 30 min at room temperature in dark. The absorbance of the samples/standards was measured at 517 nm against the blank. The decrease of the absorbance of the DPPH solution indicates an increase in DPPH radical scavenging activity. The experiment was performed in triplicates. The DPPH radical scavenging activity was calculated according to the following equation:

$$\begin{aligned} \% \text{DPPH radical scavenging activity} \\ = (A_0 - A_1) / A_0 \times 100\%, \end{aligned}$$

where A_0 is the absorbance of the control reaction and A_1 is the absorbance in the presence of the sample of the tested extracts. Percentage radical activity was plotted against the corresponding antioxidant substance concentration to obtain the IC_{50} value, which is defined as the amount of antioxidant substance required to scavenge the 50% of free radicals present in the assay system. IC_{50} values are inversely proportional to the antioxidant potential.

ABTS radical cation scavenging assay

ABTS radical cation scavenging assay was carried out by the method of Re et al. [26]. 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) ($ABTS^{2+}$) radical cation was generated by adding 14 mM ABTS to 4.9 mM

ammonium persulphate and the mixture was allowed to stand for 12–16 h in the dark at room temperature. This solution was diluted to obtain an absorbance of 0.7 ± 0.05 with ethanol at 734 nm (Shimadzu UV-Vis spectrophotometer model 2450) and the same was used for further assay. To 900 µl of ABTS radical solution, added 100 µl of the different concentrations of extracts (100–1000 µg/ml) and ascorbic acid and gallic acid standards (2.0–10 µg/ml). The solution was shaken well and the absorbance was measured after 7 min. The capacity to scavenge the ABTS radical cation was calculated using the formula same as DPPH method.

In vitro antidiabetic assays

α -amylase inhibition assay

The α -amylase inhibitory activity of the methanol extracts of bark, leaves, fruit pulp and seed of the *S.caryophyllatum* was determined according to the method described by Ranilla et al. [27]. A total of 250 µL of sample/standard and 125 µL of 0.02 M sodium phosphate buffer (pH 6.9 with 6 mM NaCl) containing α -amylase solution (0.5 mg/mL) was incubated at 25 °C for 10 min. After pre-incubation, 250 µL of 1% starch solution in 0.02 M sodium phosphate buffer (pH 6.9 with 6 mM NaCl) was added to each tube at timed intervals. The reaction mixtures were then incubated at 25 °C for 10 min. The reaction was stopped with 0.5 mL of dinitrosalicylic acid colour reagent. The test tube was then incubated in a boiling water bath for 5 min and cooled to room temperature. The reaction mixture was then diluted after adding 5 mL of distilled water and absorbance was measured at 540 nm. Acarbose was used as the positive control. The α -amylase inhibitory activity was calculated as follows;

$$\% \text{Inhibition of } \alpha\text{-amylase} = (A_0 - A_1) / A_0 \times 100\%.$$

Where, A_0 is the absorbance of the control reaction and A_1 is the absorbance in the presence of the sample of the tested extracts.

α -glucosidase inhibition assay

The α -glucosidase inhibitory activity was assessed by Dong et al. [28] with slight modifications. Briefly, a volume of 60 µL of sample/acarbose standard solution and 50 µL of 0.1 M phosphate buffer (pH 6.8) containing α -glucosidase solution (0.2 U/mL) was incubated in 96 well plates at 37 °C for 20 min. After pre-incubation, 50 µL of 5 mM *p*-nitrophenyl- α -D-glucopyranoside (PNPG) solution in 0.1 M phosphate buffer (pH 6.8) was added to each well and incubated at 37 °C for another 20 min. Then the reaction was stopped by adding 160 µL of 0.2 M Na_2CO_3 into each well, and absorbance

was recorded at 405 nm by micro-plate reader and compared to a control which had 60 μ L of buffer solution in place of the extract. The α -glucosidase inhibitory activity was expressed as percentage inhibition and was calculated same as α -amylase.

Antibacterial sensitivity tests

Antibacterial activity of bark, leaves, fruit pulp and seed extracts were investigated by the disc diffusion method [29]. Antibacterial activity was carried out against six bacteria viz., *Bacillus cereus*, *B. licheniformis*, *Escherichia coli*, *Staphylococcus aureus*, *S. hominis* and *Aerococcus viridians*. Mueller-Hinton agar medium (MHA) was used for antibacterial susceptibility tests. Petri plates were prepared by pouring 20 mL of MHA medium and allowed to solidify. The bacterial suspensions were then streaked over the media surface using a sterile cotton swab to ensure confluent growth of the organism. 20 μ L of 2.5 mg /mL of bark, leaves, fruit pulp and seed extracts were impregnated on sterile discs of 6 mm size. The discs were then aseptically placed on the surface of agar plates at well-spaced intervals. The plates were incubated at 37 °C for 24 h and the observed zones of inhibition were measured. DMSO serve as a control and antibiotic discs (6.0 mm in diameter) of 10/10 μ g/mL Ampicillin/Sulbactam, 120 μ g/mL Gentamycin were also used as positive controls. Antibacterial activity was evaluated by measuring the diameter of the zone of inhibition against tested bacteria.

The minimum inhibitory concentration (MIC) values were determined for the bacterial strains which were sensitive to the bark, leaves, fruit pulp and seed extracts in disc diffusion assay. The inoculums of the bacterial strains were prepared from 12 h broth cultures and suspensions were adjusted to 0.5 McFarland standard turbidity. The bark, leaves, fruit pulp and seed extracts of *S. caryophyllatum* dissolved in 5% dimethyl sulfoxide (DMSO) were first diluted to the highest concentration (1000 μ g/mL) to be tested, and then serial twofold dilutions were made in order to obtain a concentration range from 62.5–1000 μ g/mL in 10 mL sterile test tubes containing nutrient broth. MIC values of *S. caryophyllatum* bark, leaves, fruit pulp and seed extracts against bacterial strains were determined based on a microwell dilution method. The tubes were observed after incubation of 24 h at 37 °C. After incubation, the MIC of each sample was determined by visual inspection of the tubes. The lowest concentration of the active ingredient that inhibited growth, as detected by a lack of visual turbidity was taken to be the MIC.

Larvicidal activity

The larvicidal activity of the leaf essential oil and methanol extracts of bark, leaf, fruit pulp and seed of *S.*

caryophyllatum were tested according to the WHO [30] procedure with slight modifications. Larvicidal activity was tested against two mosquito vectors namely *Aedes aegypti* (*Ae. aegypti*) and *Culex quinquefasciatus* (*Cx. quinquefasciatus*). The larvae were obtained from Centre for Research in Medical Entomology, Madurai. The essential oil was dissolved in methanol to make the stock solution of 10,000 ppm (10 μ L/1 mL). From the stock solution, concentration of 10, 50, 100, 200 and 400 ppm were prepared. The 1% stock solution of crude extracts of bark, leaf, fruit pulp and seed were prepared with methanol. From the stock solution, 500–62.5 mg/L, dilutions were prepared with methanol. Twenty early fourth instars larvae were introduced in insects breeding box containing 100 mL of water with each concentration. A control was prepared by the addition of methanol. Larvae were fed a diet of dry yeast and dog biscuits. The numbers of dead larvae were counted after 24 h of exposure and the percentage mortality was reported from the average of three replicates. The average larval mortality data were subjected to probit analysis for calculating LD₅₀ and LD₉₀ values were calculated by using the software BioStat, 2009.

$$\text{Percent mortality} = \frac{\% \text{Mortality in treated} - \% \text{Mortality in control}}{100 - \% \text{Mortality in Control}}$$

Statistical analysis

All data were reported as mean \pm standard deviation/standard errors of means of three replicates. The IC₅₀ values were calculated using the ED50plus v 1.0 programme. Statistical analyses were performed using Microsoft Excel.

Results

The quantitative estimation of total phenol, flavonoid and tannin contents of different solvent extracts of *S. caryophyllatum* bark, leaf, fruit pulp and seed were analysed. The concentration of total phenol content expressed in terms of gallic acid equivalent (GAE) (the standard curve equation: $y = 0.188x + 0.030$, $R^2 = 0.997$). The concentrations of total phenol content in bark, leaf, fruit pulp and seed extracts of *S. caryophyllatum* ranged from 0.74 to 36.1 mg GAE/g dry weight. The highest amount of total phenol was found to be in methanol extracts of leaf (36.1 mg GAE /g), bark (31.6 mg GAE /g) and seed (29.1 mg GAE /g). While ethyl acetate extracts of leaf, bark and seed were also exhibited significant amount of phenol content with 25.7, 17.8 and 22.28 mg GAE/g respectively. The lowest concentration of total phenol content was obtained in fruit pulp extracts (Table 1).

Table 1 The quantitative estimation of total phenol content of the different solvent extracts of *S. caryophyllatum* leaf, bark fruit pulp and seed

Phytoconstituents (mg/g dry weight)	Leaf	Bark	Fruit pulp	Seed
Hexane	3.95 ± 0.2	8.4 ± 0.4	0.74 ± 0.1	5.47 ± 0.14
Ethyl Acetate	25.7 ± 0.1	17.8 ± 0.2	6.27 ± 0.15	22.28 ± 0.1
Methanol	36.1 ± 0.9	31.6 ± 0.2	8.61 ± 0.2	29.1 ± 0.3
Water	6.98 ± 0.2	21.8 ± 0.1	2.65 ± 0.1	15.5 ± 0.2

Values are given as mean ± SD of triplicate experiment

The concentration of total flavonoid content was expressed in terms of (RE) rutin equivalent (the standard curve equation: $y = 0.175x - 0.029$, $R^2 = 0.994$). The concentration of flavonoid content in bark, leaf, fruit pulp and seed extracts of *S. caryophyllatum* ranged from 0.51 to 5.63 mg RE/g dry weight. The highest concentration of flavonoid level was obtained in methanol extract from the leaf (5.63 mg RE/g) followed by bark methanol (3.6 mg RE/g) and leaf water extract (3.75 mg RE/g) respectively. Fruit pulp and seed extracts of all solvent fractions showed significant amount of flavonoid content. The lowest concentration of flavonoid was found in hexane and ethyl acetate extracts of leaves and bark (Table 2).

Total tannin content in bark, leaf, fruit pulp and seed extracts were expressed in terms of (TAE) tannic acid equivalent (the standard curve equation: $y = 0.229x + 0.041$, $R^2 = 0.998$). The maximum concentration of total tannin content was found to be 36.6 mg TAE/g in methanol extract of leaf followed by methanol bark extract (31.3 mg TAE/g), water leaf extract (26.3 mg TAE/g), ethyl acetate leaf extract (25.3 mg TAE/g) and 22.9 mg TAE/g in methanol extract of seed. Among the four plant parts tested, lowest tannin content was obtained in fruit pulp extracts (Table 3). The results of quantitative phytochemical analysis revealed that the presence of greater amounts of phytoconstituents such as phenol, flavonoids and tannins in the methanol extracts leaf, bark, fruit pulp and seed of *S. caryophyllatum* when compared to other solvent extracts.

Table 2 The quantitative estimation of total flavonoid content of the different solvent extracts of *S. caryophyllatum* leaf, bark fruit pulp and seed

Phytoconstituents (mg/g dry weight)	Leaf	Bark	Fruit pulp	Seed
Hexane	0.77 ± 0.03	0.82 ± 0.02	1.86 ± 0.03	2.03 ± 0.02
Ethyl Acetate	1.91 ± 0.01	0.51 ± 0.01	2.37 ± 0.03	2.87 ± 0.1
Methanol	5.63 ± 0.08	3.6 ± 0.06	2.5 ± 0.1	2.95 ± 0.32
Water	3.75 ± 0.06	3.3 ± 0.05	2.2 ± 0.02	2.4 ± 0.02

Values are given as mean ± SD of triplicate experiment

Table 3 The quantitative estimation of total tannin content of the different solvent extracts of *S. caryophyllatum* leaf, bark fruit pulp and seed

Phytoconstituents (mg/g dry weight)	Leaf	Bark	Fruit pulp	Seed
Hexane	15.2 ± 0.5	10.8 ± 0.1	—	3.6 ± 0.1
Ethyl Acetate	25.3 ± 0.8	16.3 ± 0.03	0.21 ± 0.08	14.1 ± 0.1
Methanol	36.6 ± 0.01	31.3 ± 0.05	0.84 ± 0.05	22.9 ± 0.1
Water	26.3 ± 0.1	16.5 ± 0.1	1.04 ± 0.1	5.1 ± 0.1

Values are given as mean ± SD of triplicate experiment

Antibacterial sensitivity test

The antibacterial activity of crude methanol extracts of bark, leaves, fruit pulp and seed was determined by disc diffusion and minimum inhibitory concentration methods against six bacterial species viz. *Bacillus cereus*, *B. licheniformis*, *Staphylococcus aureus*, *S. hominis*, *Aerococcus viridians* and *Escherichia coli*. Methanol bark extract exhibited the pronounced antibacterial activity against *S. hominis*, *B. licheniformis* and *A. viridian* with zone diameter of 12.3 mm, 12 mm and 11.3 mm respectively. Leaf extract showed potent antibacterial activity against *E. coli* (14.3 mm), *B. cereus* (14 mm), *S. aureus* (12.3 mm) but it did not exhibited any zone of inhibition against *S. hominis*. Methanol seed extract exhibited the pronounced antibacterial activity against *A. viridian*, *B. cereus* and *S. aureus* with zone diameter of 12 mm, 13 mm and 10 mm respectively (Table 4). However, fruit pulp extract exhibited moderate effect of antibacterial activity against some of the tested bacterial strains. The maximum zone of inhibition showed against *S. aureus* with zone diameter of 9 mm. The blind control (DMSO) did not inhibit any of the bacteria tested. The MIC values confirmed the significant activity against the tested microorganisms, as shown in Table 5.

Larvicidal activity

Results obtained in the present investigation on the leaf essential oil and methanol crude extracts of bark, leaf, fruit pulp and seed of *S. caryophyllatum* exhibited potent larvicidal activity in terms of LD₅₀ and LD₉₀ against the two selected mosquito vectors *Aedes aegypti* and *Culex quinquefasciatus* are presented in Table 7. The leaf essential oil of *S. caryophyllatum* showed 100% mortality against fourth instar larvae of *Ae. aegypti* and *Cx. Quinquefasciatus* at concentration of 400 ppm. At 200 ppm, over 95% mortality of fourth instar larvae of *Ae. aegypti* and 81% mortality of *Cx. Quinquefasciatus* were observed. The LD₅₀ values of the leaf essential oil recorded after 24 h of exposure were 54.8 ppm and 95.26 ppm against *Ae. aegypti* and *Cx. quinquefasciatus* respectively. The larvicidal activity increased with an increase in the leaf essential oil concentration. Similar trends were observed in the methanol crude extracts of

Table 4 Antibacterial activities of leaf, bark, fruit pulp and seed methanol extracts of *S. caryophyllatum*

Bacterial strains	Leaf	Bark	Fruit pulp	Seed	Gentamycin	Amp/Sulb
<i>S. aureus</i>	12.3 ± 0.5	8.3 ± 0.5	9 ± 0.57	10 ± 0.76	21 ± 1.1	13.5 ± 1.5
<i>S.hominis</i>	nd	12.30.5	nd	nd	28 ± 0.5	30 ± 0.5
<i>A. viridans</i>	6.3 ± 0.57	11.3 ± 0.5	6.8 ± 0.28	12 ± 0.57	24.5 ± 0.5	10 ± 1.5
<i>E. coli</i>	14.3 ± 0.5	8.3 ± 0.5	9.83 ± 0.2	7.5 ± 0.5	25 ± 0.5	10 ± 2.0
<i>B. cereus</i>	14 ± 1.0	8.4 ± 0.5	8.6 ± 0.57	13 ± 1.15	27 ± 1.1	9.5 ± 1.1
<i>B. licheniformis</i>	11.1 ± 0.5	12.1 ± 0.7	7 ± 0.57	8 ± 1.1	23 ± 1.0	31 ± 1.1

Values are given as mean ± SD of triplicate experiment

Diameter of inhibition zones (mm) including disc size 6 mm

Standard antibiotics: Ampicillin/Sulbactam (10/10 mcg/disc); Gentamycin (120 mcg/disc)

nd not detected

bark, leaf, fruit pulp and seed; and the results of mean percentage mortality against two selected mosquito vectors in different concentrations were also shown in Table 6. The methanol extracts of all plant parts exhibited lethal effects on *Ae. aegypti* and *Cx. quinquefasciatus* after 24 h of exposure at different concentrations. The highest mortality rate was found to be in methanol extract of leaf compared with other plant parts. The methanol extracts of leaf, bark and seed extracts showed over 80% mortality at 500 mg/L. Leaf methanol extract of *S. caryophyllatum* was effective against the fourth instar larvae of *Ae. aegypti* and *Cx. quinquefasciatus* with LD₅₀ values of 80.4 (LD₉₀ = 346.5 mg/L) and 128.1 mg/L (LD₉₀ = 363.9 mg/L) respectively. The bark and seed methanol extracts had significant larvicidal effect with LD₅₀ and LD₉₀ values found to be 303.6 (LD₉₀ = 541.3 mg/L) and 286.0 (LD₉₀ = 554.8 mg/L) against *Ae. aegypti* and 286.4 (LD₉₀ = 509.5 mg/L) and 300.2 (LD₉₀ = 547.2 mg/L) against *Cx. quinquefasciatus*. Methanol fruit pulp showed the least larvicidal activity compared with other three plant parts studied. At 500 mg/l concentration, it caused 71.6% and 48.3% larval mortality against *Ae.aegypti* and *Cx. quinquefasciatus* with LD₅₀ and LD₉₀ values of 387.0 (LD₉₀ = 651.6 mg/L) and 519 mg/l (LD₉₀ = 876.1 mg/L) respectively. The LD₅₀ and its lower, upper confident levels, along with the regression analyses are given in Table 7. The results obtained in this study clearly proved that the *S.*

caryophyllatum leaf essential oil and effective constituents of methanol extract of different plant parts could be considered as a potent source for the production of eco-friendly larvicidal agent against the disease causing common mosquito vectors.

Free radical scavenging assays

DPPH is a stable free radical, which has been widely accepted as a tool for estimating free radical-scavenging

Table 6 Larvicidal activity indifferent concentrations of *S. caryophyllatum* plant extracts and essential oil against *Ae. aegypti* and *Cx. quinquefasciatus* (mg/l)

Plant parts used	Concentrations (mg/l)	24 h % Mortality (<i>Ae. aegypti</i>)	24 h % Mortality (<i>Cx. Quinquefasciatus</i>)
Bark	500	83.3 ± 2	86.6 ± 5.7
	250	56.7 ± 2.8	57 ± 3
	125	15 ± 0.0	18.3 ± 2.8
Leaf	62.5	0	6.6 ± 2.8
	500	98.3 ± 2.8	96.6 ± 2.9
	250	91.7 ± 5.7	85 ± 5
Fruit pulp	125	68.3 ± 7.6	55 ± 5
	62.5	35 ± 5	26.6 ± 2.8
	500	71.6 ± 3	48.3 ± 7.6
Seed	250	28.3 ± 5.7	21.6 ± 3
	125	16.6 ± 2.8	11.6 ± 2.8
	62.5	0	0
Essential oil (ppm)	500	81.6 ± 3	81.7 ± 2.8
	250	55 ± 5	56.6 ± 3
	125	26.6 ± 2.8	15.0 ± 0.0
	62.5	0	6.6 ± 3
	400	100 ± 0.0	100 ± 0.0
	200	95 ± 5.0	81.6 ± 2.8
	100	88.3 ± 2.8	71.6 ± 3
	50	51.6 ± 3	36.7 ± 2.8
	10	18.3 ± 2.8	8.3 ± 2

Values are given as mean ± SD of triplicate experiment

Table 5 Minimum inhibitory concentration (MIC) of leaf, bark, fruit pulp and seed methanol extracts against pathogenic bacteria

Bacterial strains	MIC leaf	MIC bark	MIC Fruit	MIC Seed
<i>S. aureus</i>	500	1000	500	500
<i>S.hominis</i>	nd	500	nd	nd
<i>A. viridans</i>	nd	500	1000	250
<i>E. coli</i>	100	1000	500	1000
<i>B. cereus</i>	100	1000	500	250
<i>B. licheniformis</i>	500	500	1000	1000

nd not detected; MIC values are given in µg/ml concentration

Table 7 LD50 and other statistical analysis of *S. caryophyllatum* plant extracts and essential oil against fourth instar larvae of *Ae. aegypti* and *Cx. quinquefasciatus*

Parts used	Species	LD ₅₀ (ppm)	UCL-LCL	χ ² (df = 2)	p-level
Bark	<i>A. aegypti</i>	303.6 ± 33.8	235.9–371	0.2508	0.8821
	<i>Cx. Quinquefasciatus</i>	286.4 ± 31.7	222.8–350	0.3433	0.8423
Leaf	<i>A. Aegypti</i>	80.4 ± 37.9	4.52–156.7	0.0402	0.9801
	<i>Cx. Quinquefasciatus</i>	128.1 ± 33.5	60.87–195.3	0.0162	0.9919
Fruit pulp	<i>A. Aegypti</i>	387.3 ± 37	312–462	1.1505	0.5626
	<i>Cx. Quinquefasciatus</i>	519 ± 50.7	418–621	0.3212	0.8516
Seed	<i>A. Aegypti</i>	286 ± 38.2	209–362.8	0.7243	0.6962
	<i>Cx. Quinquefasciatus</i>	300.2 ± 35	229–370	0.3229	0.8509
Es. Oil	<i>A. Aegypti</i>	54.84 ± 20.5	13.64–96.05	0.7145	0.8698
	<i>Cx. Quinquefasciatus</i>	95.26 ± 21.2	52.87–137.6	0.4343	0.9331

Values are given as mean ± SE of triplicate experiment

activities of antioxidants. The percentage inhibition of DPPH and ABTS in the methanol extracts of bark, leaves, fruit pulp and seeds at different concentrations are shown in Figs. 1 and 2. The method is based on the reduction of methanolic-DPPH solution because of the presence of antioxidant substances having hydrogen donating groups (RH) such as phenolics and flavonoids compounds due to the formation of non radical DPPH-H form. Free radical scavenging activity indicated that methanol bark, leaves, fruit and seeds extracts have significant free radical scavenging ability on DPPH with percentage inhibition of 88.15%, 81.31%, 75.24% and 83.36% respectively. The four extracts also revealed scavenging effects on ABTS with percentage inhibition of 88.15%, 81.31%, 75.24% and 83.36% respectively. The IC₅₀ values of DPPH and ABTS radical activities of the methanol extracts of bark, leaves, fruit pulp, seeds and positive controls were shown in Table 8. A lower IC₅₀ value indicates higher antioxidant activity. The results revealed that free radical scavenging activity of methanol fruit and leaves extracts has less radical scavenging ability on DPPH with IC₅₀ value of 69.41 µg/mL, while the bark and seeds methanol extracts showed IC₅₀ value of 48.37 µg/mL compared to the positive control ascorbic acid (IC₅₀ = 9.55 µg/mL) and gallic acid (IC₅₀ = 7.05 µg/mL). The ABTS radical activities of the methanol extracts of bark, leaves, fruit pulp and seed showed IC₅₀ values of 10.26 µg/mL, 13.2 µg/mL, 69.4 µg/mL and 20.1 µg/mL respectively. The positive control ascorbic acid and gallic acid showed the IC₅₀ values of 5.6 µg/mL and 4.9 µg/mL respectively.

In vitro α-amylase and α-glucosidase assays

The in vitro α-amylase and α-glucosidase inhibitory studies of the methanol crude extracts of four different plant parts of *S. caryophyllatum* (Bark, leaf fruit pulp

and seed) showed good inhibitory activities in concentration dependent manner. Percentage inhibition was tested with different concentrations of four methanol extracts 100-500 µg/ml for α-amylase and 20-100 µg/ml for α-glucosidase. The bark extract indicated a potent inhibitory activity of α-amylase followed by seed, leaf and fruit extracts. The maximum percentage inhibitory activity of 78.03% was showed at concentration of 500 µg/ml seed extracts followed by bark (78.03%), leaf (69.4%) and fruit pulp (56.9%) at the same concentration. While the percentage inhibitory activity of four extracts also showed potent inhibition of α-glucosidase. The results showed that maximum inhibition exhibited at the concentration of 100 µg/ml by bark extract (80.9%) compared with other extracts, leaf (78.2%), seed (77.59%) and fruit pulp (63.35%). The percentage inhibition and IC₅₀ values obtained in the analyses were shown in Fig. 3a, b and Table 8.

Discussion

Phytochemical screening provides fundamental information about the medicinal importance of the plant extracts. Phenolic compounds play an important role in plant defence mechanisms against invading bacteria and other types of environmental stress, such as wounding and excessive light or ultraviolet radiation [31, 32]. These compounds can act as antioxidants by chelating metal ions, preventing radical formation and improving the antioxidant endogenous system [33]. Antioxidants which predominantly originate from phytochemicals have been reported to play an important role in the treatment of diabetes [34]. Our results also in accordance with previous report that showed high phenol content in *S. aqueum* leaf ethanol extracts may be the reason for its high anti-glycation activity [35].




Fig. 1 DPPH free radical scavenging activity of *S. caryophyllatum* **a** Ascorbic acid; **b** Gallic acid; **c** Bark methanol; **d** Leaf methanol; **e** Fruit pulp methanol; **f** Seed methanol

Syzygium species are known to be very rich in flavonoids, tannins and other polyphenol derivatives [36–39]. Flavonoids, tannins, anthocyanins and other phenolic constituents present in bark, leaves and fruits are potential antioxidants. Flavonoids are known to regenerate the damaged pancreatic β -cells in the alloxan induced diabetic rats [40]. Tannins are considered nutritionally undesirable because they precipitate proteins, inhibit digestive enzymes and affect the absorption of minerals and vitamins. Tannins were reported to exhibit antiviral, antibacterial and anti-tumour activities. However, some kinds of tannins can reduce the mutagenicity of a number of mutagens and also exhibited antioxidant, anticarcinogenic and antimicrobial activities [41].

Food rich in antioxidants plays a crucial role in the prevention of various types of cancers, cardiovascular and Alzheimer’s diseases [42]. Antioxidants which predominantly originate from phytochemicals like flavonoids and phenols have been reported to play an

important role in the treatment of diabetes [34] (Kim et al., 2011). Total antioxidant capacity may better reflect the health beneficial quality of foods than individual measurements due to the possible cooperative action of antioxidants [43]. Seeds of *S. cumini* one of the close related species of *S. caryophyllatum* have been reported to be rich in flavonoids, which account for the scavenging of free radicals and a protective effect on antioxidant enzymes and they have also been found to have high total phenolic content with significant antioxidant activity [44, 45]. The results obtained in present study are in agreement with previous studies which showed that high total phenolic content increases the antioxidant activity [46]. According to previous reports the close related species *S. cumini*, *S. aromaticum* fruits and leaves extract have been reported to have high antioxidant activity [47, 48].

Different parts of *S. cumini* such as seeds, bark, fruit and leaves have been used in traditional medicine as a




Fig. 2 ABTS free radical scavenging activity of *S.caryophyllatum* **a** Ascorbic acid; **b** Gallic acid; **c** Bark methanol; **d** Leaf methanol; **e** Fruit pulp methanol; **f** Seed methanol


Table 8 Free radical scavenging activities of *S. caryophyllatum* bark, leaf, fruit pulp, seed methanol extracts and standard antioxidants (Ascorbic acid and Gallic acid)

Samples	IC50 (µg/ml)			
	DPPH	ABTS	α -amylase	α -glucosidase
Bark	19.07 ± 0.2	10.26 ± 0.06	191 ± 3.2	53.75 ± 1.6
Leaf	34.9 ± 1.2	13.2 ± 0.03	265.4 ± 5.7	57.8 ± 0.3
Fruit pulp	69.4 ± 0.7	120.2 ± 0.4	291.5 ± 4.3	61.8 ± 0.1
Seed	41.68 ± 0.6	20.1 ± 0.14	265 ± 1.8	56.1 ± 0.2
Ascorbic acid	9.55 ± 0.01	5.6 ± 0.03	–	–
Gallic acid	7.05 ± 0.09	4.9 ± 0.05	–	–
Acarbose	–	–	194.2 ± 5.1	41.13 ± 0.4

Each values in the table was obtained by calculating the average of three analysis ± SE

remedy for diabetes mellitus in many countries [49, 50]. The use of alpha-amylase and alpha-glucosidase enzyme inhibitors could be a prospective therapeutic agent for the effective management of postprandial hyperglycemia in diabetes mellitus patients [51]. It is obvious that flavonoids and polyphenols have been shown to inhibit pancreatic alpha-amylase and intestinal alpha-glucosidase enzymes and also reported that there is a positive correlation between the inhibition of alpha-amylase and alpha-glucosidase and flavonoids and polyphenol content [52–54]. Our investigation of in vitro studies of bark, leaf, fruit pulp and seed methanol extracts of *S. caryophyllatum* also showed remarkable inhibition of alpha-amylase and alpha-glucosidase suggesting the presence of potential enzyme inhibiting compounds in the plant extracts.

The use of plant extract for mosquito control save the environment from synthetic chemicals. Plant products are emerging as a potential source of mosquito control



and among them essential oil have special interest due to their insecticidal properties [55, 56]. In general essential oils from plant sources have been considered important natural resources to act as insecticides [57, 58]. Essential oils derived from various plants not only exhibit inhibitory activity against fungi, bacteria and dermatites but also show strong mosquito larvicidal and repellent activities [59]. The finding of the present study compare well with previous finding of Bagavan and Rahuman [60] who reported that the hexane crude extract of *S. aromaticum* flower bud exhibited potent larvicidal activity against fourth instar of *Anopheles vagus* and *Cx. vishnui*. A similar study conducted by Fayemiwo et al. [61] reported that the flower bud oil of *S. aromaticum* was effective against the *Ae. aegypti* ($LC_{50} = 92.56$ mg/l) and *Cx. quinquefasciatus* ($LC_{50} = 124.4$ mg/l) larvae. Our results are also comparable to earlier reports of Kaushik and Saini [62] who observed that the highest mortality was found in methanol leaf extract of *S. cumini* against the larvae of *Ae. aegypti*. The results of the present study suggested that the larvicidal potential of leaf essential oil and plant extracts of *S. caryophyllatum* and emphasize need to incorporate them in vector control and management programmes.

Conclusion

Based on the results of bioprospecting and biological screening of *S. caryophyllatum* proved to be a good medicinal plant used for curing various ailments in different traditional systems of medicine. The results of bioprospecting investigation the various polyphenolic compounds present in the plant extracts attribute the antioxidant, alpha amylase, alpha glucosidase, antibacterial and larvicidal activities. Thus the present study suggests that *S. caryophyllatum* bark, leaf, fruit pulp and seed can be used as natural antioxidant source to prevent diseases associated with free radicals. Also, this plant can be a good source for further purification

studies for isolation and characterization of compounds related to these antioxidants, antidiabetic and antibacterial activities. These findings justify and confirm the traditional uses of this plant. Further research is necessary in order to know all the active principles and their pharmacological properties.

Abbreviations

ABTS: -2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid); DMSO: Dimethyl sulfoxide; DPPH: 1,1-diphenyl-2-picrylhydrazyl; GAE: Gallic acid equivalent; IC_{50} : 50% Inhibitory concentration; LD_{50} : 50% Lethal Dose; MHA: Mueller-Hinton agar medium; mM: Milli molar; Na_2CO_3 : Sodium Carbonate; NaCl: Sodium Chloride; PNPG: *p*-nitrophenyl- α -D-glucopyranoside; RE: Rutin equivalent; TAE: Tannic acid equivalent

Acknowledgments

The authors would like to express their gratitude to Centre for Research in Medical Entomology (CRME), Madurai, India for providing mosquitos eggs for larvicidal analysis.

Funding

First author thanks University Grant Commission for the award of meritorious fellowship. Corresponding author thanks partial financial support through University Grant Commission-Centre of Advanced Study (UGC-CAS), Department of Biotechnology- Interdisciplinary Programme of Life Sciences for Advanced Research (DBT-IPLS) and Department of Science and Technology (DST)-Purse programmes.

Availability of data and materials

Not Applicable.

Authors' contributions

Both the authors are equally contributed. Both the authors read and approved the final manuscript.

Authors' information

Dr. Stalin: Completed his PhD in Department of Plant Science, School of Biological Sciences, Madurai Kamaraj University, Madurai. His area of research is species distribution modelling, bioprospecting and conservation of selected medicinal plant through phenological studies and in vitro propagation.

Dr. P. Sudhakar Swamy: Professor, Department of Plant Science, School of Biological Sciences, Madurai Kamaraj University, Madurai. His area of specialisation is Forest ecology, invasion ecology, bioprospecting of medicinal plants and their conservation.

Ethics approval and consent to participate

Not Applicable.

Consent for publication

Not Applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 30 October 2017 Accepted: 27 December 2017

Published online: 18 January 2018

References

- Gupta SK. Pharmacology and therapeutics in the new millennium. New Delhi: Narosa Publishing House; 2001.
- Wong C, Li H, Cheng K, Chen F, et al. A systematic survey of antioxidant activity of 30 Chinese medicinal plants using the ferric reducing antioxidant power assay. *Food Chem.* 2006;97:705–11.
- Ames SN, Shigenaga MK, Hagen TM, et al. Oxidants, antioxidant and the degenerative diseases of aging. *Proc Natl Acad Sci U S A.* 1993; 90:7915–22.
- Shukla VKS, Wanasundara PKJP, Shahidi F, et al. Natural antioxidants from oilseeds. In: Shahidi F, editor. *Natural antioxidants: chemistry, health effects, and applications.* Champaign: AOCS Press; 1997. p. 97–132.
- Mohamed AA, Khalil AA, El-Beltagi HES, et al. Antioxidant and antimicrobial properties of kaff maryam (*Anastatica hierochuntica*) and doum palm (*Hyphaene thebaica*). *Grasas Y Aceites.* 2010;61:67–75.
- Ozsoy N, Can A, Yanardag R, Akev A, et al. Antioxidant activity of *Smilax excelsa* leaf extracts. *Food Chem.* 2010;X110:571–83.
- Giovanelli G, Buratti S. Comparison of polyphenolic composition and antioxidant activity of wild Italian blueberries and some cultivated varieties. *Food Chem.* 2009;112:903–8.
- Salah N, Miller NJ, Paganga G, Tijburg L, Bolwell GP, et al. Polyphenolic flavanols as scavengers of aqueous phase radicals and as chain breaking antioxidants. *Arch Biochem Biophys.* 1995;322:339–46.
- Oomah BD, Cardador-Martinez A, Loarca-Pina G, et al. Phenolics and antioxidative activities in common beans. *J Sci Food Agric.* 1995; 85:935–42.
- Brighente IMC, Dias M, Verdi LG, Pizzolatti MG. Antioxidant activity and total phenolic content of some Brazilian species. *Pharm. Biol.* 2007;45:156–61.
- Salazar R, Pozes ME, Cordero P, Perez J, Salinas MC, Waksman N. Determination of the antioxidant activity of plants from Northeast Mexico. *Pharm Biol.* 2008;46:166–70.
- Ahmad I, Aqil F. Vitro efficacy of bioactive extracts of 15 medicinal plants against SBL-producing multidrugresistant enteric bacteria. *Microbiol Res.* 2007;162:264–75.
- Barbour EK, Al Sharif M, Sagherian VK, Habre AN, Talhouk RS, Talhouk SN. Screening of selected indigenous plants of Lebanon for antimicrobial activity. *J Ethnopharmacol.* 2004;93:1–7.
- Iwu MW, Duncan AR, Okunji CO. New antimicrobials of plant origin. In: Janick J, editor. *Perspectives on new Crops and New Uses.* Alexandria: ASHS Press; 1999. p. 457–62.
- WHO (2013). *World Filariasis Report.* Geneva: Switzerland; 282.
- WHO (2015). *Dengue and Severe Dengue Fact Sheet.* World Malaria Report. Washington, DC: National Press Club; 1–5.
- Ediriweera ERHSS, Ratnasooriya WDA. Review on herbs used in treatment of diabetes mellitus by Sri Lankan and traditional physicians. *Ayu.* 2009;30:373–91.
- Shilpa KJ, Krishnakumar G. Phytochemical screening, antibacterial and antioxidant efficacy of the leaf and bark extracts of *Syzygium caryophyllatum* (L.) Alston. *Int J Pharm Pharm Sci.* 2012;4:198–202.
- Rabeque CS, Padmavathy S. Comparative phytochemical analysis of root extracts of *S. caryophyllatum* (L.) Alston and *S. densiflorum* wall. *Pharmacie Globale (IJCP).* 2014;04:1–3.
- Harsha VH, Sripathy V, Hedge GK. Ethnoveterinary practices in Uttara Kannada district of Karnataka. *Indian J Tradit Knowl.* 2005;4:253–8.
- Ratheesh Narayanan MK, Anilkumar N, Balakrishnan V, Sivadasan M, Ahmed Alfarhan H, Alatar AA. Wild edible plants used by the Kattunaikka, Paniya and Kuruma tribes of Wayanad District, Kerala, India. *J Med Plant Res.* 2011;5:3520–9.
- Lister B, Wilson P. Measurement of total phenolics and ABTS assay for antioxidant activity (personal communication). Crop Research Institute: Lincoln; 2001.
- Quettier DC, Gressier B, Vasseur J, Dine T, Brunet C, Luyckx MC, Cayin JC, Bailleul F, Trotin F. Phenolic compounds and antioxidant activities of buckwheat (*Fagopyrum esculentum* Moench) hulls and flour. *J Ethnopharmacol.* 2000;72:35–42.
- Shanmugam T, Sathish Kumar T, Panneer Selvam K. *Laboratory handbook on biochemistry.* New Delhi: PHI Learning Pvt. Ltd.; 2010.
- Shimada K, Fujikawa K, Yahara K, Nakamura T. Antioxidative properties of xanthan on the autoxidation of soybean oil in cyclodextrin emulsion. *J Agric Food Chem.* 1992;40:945–8.
- Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med.* 1999;26:1231–7.
- Ranilla LG, Kwon YI, Genoves MI, Lajolo FM, Shetty K. Antidiabetes and antihypertension potential of commonly consumed carbohydrate sweeteners using *in vitro* models. *J Med Food.* 2008;11(2):337–48.
- Dong HQ, Li M, Zhu F, Liu FL, Huang JB. Inhibitory potential of trilobatin from *Lithocarpus polystachyus* Rehd against α -glucosidase and α -amylase linked to type 2 diabetes. *Food Chem.* 2012;130:261–6.
- Irobi ON, Moo-Young M, Anderson WA, Daramola SO. Antimicrobial activity of the bark of *Bridelia ferruginea* (Euphorbiaceae). *Int J Pharmacog.* 1994;34:87–90.
- World Health Organization. Report of the WHO informal consultation on the evaluation on the testing of insecticides, CTD/WHO PES/IC/96.1. Geneva: WHO; 1996. p. 69.
- Harbourne JB. *The flavonoids: advances in research since 1986.* London: Chapman & Hall; 1994.
- Wallace G, Fry SC. Phenolic components of the plant cell wall. *Int Rev Cytol.* 1994;151:229–67.
- Al-Azzawie HF, Mohamed-Saïel SA. Hypoglycemic and antioxidant effect of oleuropein in alloxan-diabetic rabbits. *Life. Science.* 2006;78:1371–7.
- Kim JS, Hyun TK, Kim MJ. The inhibitory effects of ethanol extracts from *Sorghum*, foxtail millet and proso millet on α -glucosidase and α -amylase activities. *Food Chem.* 2011;124:1647–51.
- Palanisamy UD, Ling LT, Manoharan T, Sivapalan V, Subramaniam T, Helme MH. Standardized extract of *Syzygium aqueum*: a safe cosmetic ingredient. *Int J Cosmet Sci.* 2011;1:7.
- Chattopadhyay D, Sinha BK, Vaid LK. Antibacterial activity of *Syzygium* species. *Fitoterapia.* 1998;69:356–67.
- Mahmoud II, Marzouk MSA, Moharram FA, El-Gindi MR, Hassan AMK. Acylated flavonol glycosides from *Eugenia jambolana* leaves. *Phytochemistry.* 2001;58:1239–44.
- Sharma SB, Nasir A, Prabhu KM, Dev G, Murthy PS. Hypoglycemic and hypolipidemic effect of ethanolic extracts of seeds of *E. Jambolana* in alloxan-induced diabetic model of rabbits. *J Ethnopharmacol.* 2003;85:201–6.
- Reynertson KA, Yang H, Jiang B, Basile MJ, Kennelly MEJ. Quantitative analysis of antiradical phenolic constituents from fourteen edible Myrtaceae fruits. *Food Chem.* 2008;109:883–90.
- Hakkim FL, Giriya S, Kumar RS, Jalaluddeen MD. Effect of aqueous and ethanolic extracts of *Cassia auriculata* L. flowers on diabetes using alloxan induced diabetic rats. *Int J diabetes and Metabolism.* 2007;15:100–6.
- Chung KT, Wei C, Johnson MG. Are tannins a double edged sword in biology and health. *Trends Food Sci Technol.* 1998;9:168–75.
- Cui K, Luo XL, KY X, Murthy MRV. Role of oxidative stress in neurodegeneration: recent developments in assay methods for oxidative stress and nutraceutical antioxidants. *Prog Neuro-Psychopharmacol Biol Psychiatry.* 2004;28:771–99.
- Ghiselli A, Serafini M, Natella F, Scaccini C. Total antioxidant capacity as a tool to assess redox status: critical view and experimental data. *Free Rad Biol Med.* 2000;29:1106–14.
- Ravi K, Ramachandran B, Subramanian S. Protective effect of *Eugenia jambolana* seed kernel on tissue antioxidants in streptozotocin induced diabetic rats. *Biol Pharm Bull.* 2004;27:1212–7.
- Bajpai M, Pande A, Tewari SK, Prakash D. Phenolic contents and antioxidant activity of some food and medicinal plants. *Int J Food Sci Nutr.* 2005;56:287–29.

46. Abhishek KS, Vinod KV. *Syzygium cumini*: An overview. *J Chem Pharm Res*. 2011;3:108–11.
47. Ruan ZP, Zhan LL, Lin YM. Evaluation of the antioxidant activity of *Syzygium cumini* leaves. *Molecules*. 2008;13:2545–56.
48. Jirovetz L, Buchbauer G, Stoilova I, Stoyanova A, Krastanov A, Schmidt E. Chemical composition and antioxidant properties of clove leaf essential oil. *J Agric Food Chem*. 2006;54:6303–7.
49. Indira G, Mohan R. Jamun Fruits, National Institute of Nutrition. Hyderabad: Indian Council of Medical Research; 1993. p. 34–7.
50. Teixeira CC, Pinto LP, Kessle FHP, Knijnik L, Pinto CP, Gastaldo GJ, Fuchs FD. The effect of *Syzygium cumini* (L.) Skeels on post-prandial blood glucose levels in non-diabetic rats and rats with streptozotocin-induced diabetes mellitus. *J Ethnopharmacol*. 1997;56:209–13.
51. Ali H, Houghton PJ, Soumyanath A. α -amylase inhibitory activity of some Malaysian plants used to treat diabetes; with particular reference to *Phyllanthus amarus*. *J Ethnopharmacol*. 2006;107:449–55.
52. Adisakwattana S, Ruengsamran T, Kampa P, Sompong W. In vitro inhibitory effects of plant based foods and their combinations on intestinal α -glucosidase and pancreatic α -amylase. *BMC Complement Altern Med*. 2012;12:110.
53. Pereira DF, Cazarolli LH, Lavado C, Mengatto V, Figueiredo MS, Guedes A, Pizzolatti MG, Silva FR. Effect of flavonoids on α -glucosidase activity: potential targets for glucose homeostasis. *Nutrition*. 2011;27:1161–7.
54. Xiao J, Kai G, Ni X, Yang F, Chen X. Interaction of natural polyphenols with α -amylase in vitro: molecular property-affinity relationship aspect. *Mol Biosyst*. 2011;7:1883–90.
55. Cheng SS, Chang HT, Chang ST, Tsai KH, Chen WJ. Bioactivity of selected plant essential oils against yellow fever mosquitoes *Aedes aegypti* larvae. *Bioresour Technol*. 2003;89:99–102.
56. Sukumar K, Perich MJ, Boobar LR. Botanical derivatives in mosquito control: a review. *J Am Mosq Control Assoc*. 1991;7:210–37.
57. Gbolade AA. Plant-derived insecticides in the control of malaria vector. *J Trop Med Plants*. 2001;2:91–7.
58. Adebayo TA, Gbolade AA, Olaifa JJ. Comparative study of toxicity of essential oils to larvae of three mosquito species. *Nig J Nat Med*. 1999;3:74–6.
59. Cheng SS, Liu JY, Tsai KH, Chen WJ, Chang ST. Chemical composition and mosquito larvicidal activity of essential oils from leaves of different *Cinnamomum osmophloem* provenances. *J Agric Food Chem*. 2004;52:4395–400.
60. Bagavan A, Rahuman AA. Evaluation of larvicidal activity of medicinal plant extracts against three mosquito vectors. *Asian Pac J Trop Med*. 2011;4:29–34.
61. Fayemiwo KA, Adeleke MA, Okoro OP, Awojide SH, Awoniyi IO. Larvicidal efficacies and chemical composition of essential oils of *Pinus sylvestris* and *Syzygium aromaticum* against mosquitoes. *Asian Pac J Trop Biomed*. 2014;4(1):30–4.
62. Kaushik R, Saini P. Screening of some semi-arid region plants for larvicidal activity against *Aedes aegypti* mosquitoes. *J Vector Borne Dis*. 2009;46:244–6.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
